

**Ecological Vegetation Class Coverage: Pre-European Settlement and 1993<sup>1</sup>**

		Ecological Vegetation Classes, complexes and mosaics.		Pre-European <sup>^</sup>		Vegetation Cover (1993)				
				Total Area	Total Extant Area	Private (Freehold) Land		Conservation Reserves (includes Forests SPZ)	Other Public Land (includes unknown)	
Bioregion	No.	Conservation Status	Ha	Ha	%*	Ha	% of 1993 Cover that is Freehold	Ha	Ha	
Murray Fans	22	Endangered	187,992	5313	3%	2403	45%	540	2369	
	2	Vulnerable	6559	456	7%	56	12%	329	71	
	2	Depleted	67,829	39,144	58%	2509	6%	6360	30,275	
	1	Least Concern	2295	2039	89%	0	0%	1333	706	
<b>TOTAL</b>	<b>27</b>		<b>264,676</b>	<b>46,951</b>	<b>18%</b>	<b>4968</b>	<b>11%</b>	<b>8562</b>	<b>33,421</b>	
Victorian Riverina	3	Presumed Extinct	1482	0		0		0	1	
	42	Endangered	719,812	11,723	2%	7725	66%	1780	2219	
	9	Vulnerable	19,937	3269	16%	2271	69%	253	745	
	2	Depleted	8272	5719	69%	1534	27%	605	3581	
	1	Least Concern	212	22	10%	5	23%	15	2	
<b>TOTAL</b>			<b>749,714</b>	<b>20,733</b>	<b>2.8%</b>	<b>11,535</b>	<b>56%</b>	<b>2653</b>	<b>6548</b>	
Northern Inland Slopes	2	Presumed Extinct	821	0	-	0	-	0	0	
	19	Endangered	57,803	1342	2%	901	67%	303	138	
	2	Vulnerable	24,799	2272	9%	1248	55%	291	734	
	3	Depleted	625	418	67%	88	21%	329	0	
	3	Least Concern	8680	5951	69%	1992	33%	3364	595	
<b>TOTAL</b>			<b>92,727</b>	<b>9983</b>	<b>11%</b>	<b>4228</b>	<b>42%</b>	<b>4288</b>	<b>1467</b>	
Goldfields	24	Endangered	48,220	3158	7%	1078	34%	41	2039	
	8	Vulnerable	11,637	1640	14%	913	56%	18	709	
	5	Depleted	101,068	55,450	55%	8242	15%	1815	45,393	
	1	Least Concern	9288	5227	56%	1067	20%	1150	3010	
<b>TOTAL</b>			<b>170,213</b>	<b>65,474</b>	<b>38%</b>	<b>11,299</b>	<b>17%</b>	<b>3024</b>	<b>51,151</b>	
Central Victorian Uplands	24	Endangered	185,973	6765	4%	4522	67%	1280	963	
	13	Vulnerable	162,232	11,355	7%	9493	84%	1209	653	
	5	Depleted	153,747	63,489	41%	33,181	52%	18,674	11,634	
	1	Rare	103	27	26%	27	100%	0	0	
	5	Least Concern	20,883	10,063	48%	4262	42%	2767	3034	
<b>TOTAL</b>			<b>522,937</b>	<b>91,968</b>	<b>18%</b>	<b>51,485</b>	<b>56%</b>	<b>23,930</b>	<b>16,283</b>	
Highlands - Northern Fall	6	Endangered	4231	675	16%	624	92%	15	36	
	7	Vulnerable	10,580	4549	43%	854	19%	2757	937	
	3	Depleted	5744	2785	48%	1274	46%	715	796	
	2	Rare	581	444	76%	48	11%	391	5	
	17	Least Concern	493,938	381,002	77%	44,935	12%	117,686	218,381	

<b>TOTAL</b>			515,075	389,454	76%	47,734	12%	121,564	220,157
<b>Highlands - Southern Fall</b>	1	Vulnerable	6	6	100%	0	0%	6	0
	9	Least Concern	966	900	93%	16	2%	257	627
<b>TOTAL</b>			972	906	93%	16	2%	263	627
<b>Victorian Alps</b>	2	Vulnerable	1092	1035	95%	0	0	762	273
	2	Rare	2565	2565	100%	0	0	1276	1289
	15	Least Concern	86,727	86,419	100%	16	0	30,069	56,318
<b>TOTAL</b>			90,384	90,019	100%	16	0%	32,106	57,880
<b>TOTAL for GB Catchment</b>			<b>2,406,699</b>	<b>715,443</b>	<b>30%</b>	<b>131,250</b>	<b>18%</b>	<b>202,003</b>	<b>387,929</b>

\* percentage of original Ecological Vegetation Class remaining. NRE's 1993 Tree 100 layer used to calculate areas remaining against the bioregional boundaries updated in 2002.

Minor discrepancies occur due to different scales of data captured (1:250,00 and 1:100,000).

^Terminology used in this Goulburn Broken Native Vegetation Management Strategy is

*pre-European* rather than *pre-1750* to simplify communication. (Pre-1750 is terminology used on NRE Corporate Database from which EVC information is extracted.)

*Information derived from NRE's Corporate database.*

Footnote – some subtotals may differ when added in this table due to the rounding effect of raw data.